



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

318. Proposed by PROFESSOR R. D. CARMICHAEL, Anniston, Ala.

Sum to infinity the series $n/(4n^2 - 1)^2$ beginning with $n=1$.

319. Proposed by C. N. SCHMALL, New York City.

A man desires to purchase eggs at 5 cents, 1 cent, and $\frac{1}{2}$ cent, respectively, in such numbers that he will obtain 100 eggs for a dollar. How many solutions in rational integers?

320. Proposed by FRANCIS RUST, C. E., Pittsburg, Pa.

Solve for t , $\cos t = m \cos 2t$.

GEOMETRY.

345. Proposed by LLOYD HOLSINGER, Bradley Polytechnic Institute, Peoria, Ill.

If a variable polygon move in such a way that its n sides turn severally round n fixed points O_1, O_2, \dots, O_n while $n-1$ of its vertices slide, respectively, along $n-1$ fixed straight lines v_1, v_2, \dots, v_{n-1} , then the last vertex will describe a conic; and the locus of the point of intersection of any pair of non-adjacent sides will also be a conic. Cremona's *Projective Geometry*.

346. Proposed by G. I. HOPKINS, M. A., Professor of Mathematics and Astronomy, High School, Manchester, N. H.

Prove the theorem for finding the lateral area of a frustum of a cone without the use of the theory of limits.

347. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

ABC is a triangle, and D, E, F are the mid points of the arcs of its nine-point circle cut off by BC, CA, AB , respectively. The inscribed circle touches these sides at X, Y, Z . Are the lines DX, EY, FZ concurrent? A purely geometrical discussion required.

CALCULUS.

276. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

In a certain country the tax per \$1 on a person's income varies as the cube root of the number of dollars, and when the income is \$8000 the rate per dollar is 5 cents. Find the largest net income possible.

277. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

Find $\frac{d^2x}{ds^2}$ and $\frac{d^2y}{ds^2}$ for $y = c \sinh \frac{x}{c}$.

278. Proposed by S. A. COREY, Hiteman, Iowa.

If C be Euler's constant, .577,215,664,9... and if B_1, B_2, B_3 , etc., be Bernoulli's numbers, $\frac{1}{6}, \frac{1}{36}, \frac{1}{42}$, etc., prove that

$$C = \frac{1}{2} + \frac{B_1}{2} - \frac{B_2}{4} + \frac{B_3}{6} - \frac{B_4}{8} + \dots - (1)^m \frac{B_m}{2m} + \dots$$